


New solutions to the tetrahedron equation associated with quantized six-vertex models

Mathematical Society of Japan Spring Meeting
Chuo University@2023/03/18

Akihito Yoneyama

Institute of Physics, University of Tokyo, Komaba

Based on: A. Kuniba, S. Matsuike, AY, arXiv:2208.10258



- Matrix equation on $V_1 \otimes \dots \otimes V_6$ (V_i : linear space) [Zamolodchikov'81]
 - X_{ijk} ($X = A, B, C, D$) acts non-trivially only on $V_i \otimes V_j \otimes V_k$.

$$A_{124}B_{135}C_{236}D_{456} = D_{456}C_{236}B_{135}A_{124}$$

- 3D analog of Yang-Baxter equation (YBE)
 - We can construct a 3D version of transfer matrices similarly to YBE.
- Several solutions are known although less systematic than YBE.
 - Zamolodchikov, Baxter, Bazhanov, Korepanov, Mangazeev, Sergeev, Stroganov, Kapranov, Voevodsky, Kazhdan, Soibelman, Carter, Saito, Kuniba, Okado, ...

RLLL relation

- Today, we focus on the *RLLL* type tetrahedron equation:

$$L_{124}L_{135}L_{236}R_{456} = R_{456}L_{236}L_{135}L_{124}$$

- If we specify the outer lines for 1,2,3-th spaces, this reads as

$$\sum_{\alpha,\beta,\gamma} \left[\begin{array}{c} c \\ i \swarrow \color{orange}{5} \alpha \\ \color{orange}{\gamma} \color{orange}{6} \beta \searrow 4 \\ j \swarrow \color{orange}{6} \beta \searrow \color{orange}{4} \\ k \end{array} \right] \circ R_{456} = \sum_{\alpha,\beta,\gamma} R_{456} \circ \left[\begin{array}{c} c \\ i \swarrow \color{orange}{4} \beta \color{orange}{6} \\ \color{orange}{\gamma} \color{orange}{6} \color{orange}{\alpha} \color{orange}{5} \\ j \swarrow \color{orange}{4} \alpha \color{orange}{5} \searrow a \\ k \end{array} \right] \dots (*)$$

$$L(|i\rangle \otimes |j\rangle \otimes |k\rangle) = \sum_{a,b} |a\rangle \otimes |b\rangle \otimes L_{i,j}^{a,b} |k\rangle$$

$$\begin{array}{c} b \\ \uparrow \\ i \longrightarrow a \\ \downarrow \\ j \end{array} = L_{i,j}^{a,b}$$

- For each (i, j, k, a, b, c) , $(*)$ gives linear equations for R .
- If we can ansatz "good" L s, we can then obtain a solution to the *RLLL* type tetrahedron equation by solving these equations.
- In fact, it can be done by considering a quantized six vertex model for L s.

■ q -Oscillator algebra O_q

□ Genetators: $\mathbf{k}, \mathbf{a}^\pm$


□ Relations $\mathbf{k}\mathbf{a}^\pm = q^\pm \mathbf{a}^\pm \mathbf{k}$, $\mathbf{a}^+ \mathbf{a}^- = 1 - \mathbf{k}^2$, $\mathbf{a}^- \mathbf{a}^+ = 1 - q^2 \mathbf{k}^2$

□ Representation π_O on $F_+ = \bigoplus_{m \in \mathbb{Z}_{\geq 0}} \mathbb{C} |m\rangle$:

$$\pi_O : \mathbf{k} |m\rangle = q^m |m\rangle, \quad \mathbf{a}^+ |m\rangle = |m+1\rangle, \quad \mathbf{a}^- |m\rangle = (1 - q^2) |m-1\rangle$$

■ L -operator $L^O \in \text{End}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes F_+)$ [Bazhanov-Sergeev'06]

$$L^O(|i\rangle \otimes |j\rangle \otimes |k\rangle) = \sum_{a,b \in \{0,1\}} |a\rangle \otimes |b\rangle \otimes \pi_O((L^O)_{i,j}^{a,b} |k\rangle) \quad (L^O)_{i,j}^{a,b} = i \begin{array}{c} \uparrow b \\ \hline \rightarrow a \\ \hline \downarrow j \end{array}$$



μ : parameter

■ Thm: [Bazhanov-Sergeev'06]

□ Consider the following *RLLL* relation for L^O :

$$\underbrace{L_{124}^O}_{\mu_4} \underbrace{L_{135}^O}_{\mu_5} \underbrace{L_{236}^O}_{\mu_6} R_{456}^{OOO} = R_{456}^{OOO} L_{236}^O L_{135}^O L_{124}^O$$

□ $R^{OOO} \in \text{End}(F_+^{\otimes 3})$ is uniquely determined and given by

$$(R^{OOO})_{i,j,k}^{a,b,c} = \delta_{i+j}^{a+b} \delta_{j+k}^{b+c} \left(\frac{\mu_3}{\mu_2}\right)^i \left(-\frac{\mu_1}{\mu_3}\right)^b \left(\frac{\mu_2}{\mu_1}\right)^k q^{ik+b(k-i+1)} \binom{a+b}{a}_{q^2} {}_2\phi_1 \left(\begin{matrix} q^{-2b}, q^{-2i} \\ q^{-2a-2b} \end{matrix}; q^2, q^{-2c} \right)$$

$$(z; q)_\infty = \prod_{n \geq 0} (1 - zq^n) \quad (z; q)_m = \frac{(z; q)_\infty}{(zq^m; q)_\infty} \quad {}_2\phi_1 \left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix}; q, z \right) = \sum_{n \geq 0} \frac{(\alpha; q)_n (\beta; q)_n}{(\gamma; q)_n (q; q)_n} z^n$$

□ R^{OOO} also satisfies the *RRRR* type tetrahedron equation:

$$R_{124}^{OOO} R_{135}^{OOO} R_{236}^{OOO} R_{456}^{OOO} = R_{456}^{OOO} R_{236}^{OOO} R_{135}^{OOO} R_{124}^{OOO}$$

■ Thm: [Kapranov-Voevodsky'94]

□ R^{OOO} = intertwiner of irreps of quantum coordinate ring $A_q(A_2)$

$$R^{OOO} \circ (\pi_1 \otimes \pi_2 \otimes \pi_1(\Delta^{\text{op}}(g))) = (\pi_2 \otimes \pi_1 \otimes \pi_2(\Delta(g))) \circ R^{OOO} \quad \forall g \in A_q(A_2)$$

q -Weyl algebra

- Summary of [Bazhanov-Sergeev'06]:
 - We can obtain both $RLLL$ & $RRRR$ type tetrahedron equations.
 - Obtained R^{000} is characterized as intertwiner of $A_q(A_2)$.

- Aim: Generalization of [Bazhanov-Sergeev'06]

- Our approach: q -Oscillator algebra \rightarrow q -Weyl algebra

- Parametric generalization of L^0 by an embedding $O_q \hookrightarrow W_q$:

$$\mathbf{k} \mapsto X, \quad \mathbf{a}^+ \mapsto Z, \quad \mathbf{a}^- \mapsto Z^{-1}(1 - X^2)$$

- q -Weyl algebra W_q

- Generators: $X^{\pm 1}, Z^{\pm 1}$

- Relations: $XZ = qZX$

- Representations π_X, π_Z on $F = \bigoplus_{m \in \mathbb{Z}} \mathbb{C} |m\rangle$:

$$\pi_X : X |m\rangle = q^m |m\rangle, \quad Z |m\rangle = |m + 1\rangle \quad (\text{coordinate rep})$$

$$\pi_Z : X |m\rangle = |m - 1\rangle, \quad Z |m\rangle = q^m |m\rangle \quad (\text{momentum rep})$$

q -Weyl algebra valued six vertex model


7/15

■ L -operators L^A ($A = X, Z, O$) [Kuniba-Matsuike-Y'22]

▣ $L^A \in \text{End}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes F)$ ($A = X, Z$)

▣ $L^O \in \text{End}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes F_+)$

$$L^A(|i\rangle \otimes |j\rangle \otimes |k\rangle) = \sum_{a,b \in \{0,1\}} |a\rangle \otimes |b\rangle \otimes \pi_A((L^A)_{i,j}^{a,b} |k\rangle) \quad (A = X, Z, O)$$



$$(L^X)_{i,j}^{a,b} = (L^Z)_{i,j}^{a,b} \quad r \quad s \quad twX \quad -qtX \quad Z \quad Z^{-1}(rs - t^2wX^2)$$

$$(L^O)_{i,j}^{a,b} \quad 1 \quad 1 \quad \mu\mathbf{k} \quad -q\mu^{-1}\mathbf{k} \quad \mathbf{a}^+ \quad \mathbf{a}^-$$

r, s, t, w, μ : parameters

■ Remark:

▣ L^X for $(r, s, t, w) = (1, 1, \mu^{-1}, \mu^2)$ corresponds to L^O via the pullback.

▣ L^Z doesn't have such a correspondence and behaves differently from L^O .

Main result

■ Thm: [Kuniba-Matsuike-Y'22]

- We solved *RLL* relations for some *ABC*s.
- Each *L* has different parameters depending on its tensor component.

$$\underbrace{L_{124}^A}_{r_4, s_4, t_4, w_4 \text{ or } \mu_4} \underbrace{L_{135}^B}_{r_5, s_5, t_5, w_5 \text{ or } \mu_5} \underbrace{L_{236}^C}_{r_6, s_6, t_6, w_6 \text{ or } \mu_6} R_{456}^{ABC} = R_{456}^{ABC} \underbrace{L_{236}^C}_{r_6, s_6, t_6, w_6 \text{ or } \mu_6} \underbrace{L_{135}^B}_{r_5, s_5, t_5, w_5 \text{ or } \mu_5} \underbrace{L_{124}^A}_{r_4, s_4, t_4, w_4 \text{ or } \mu_4}$$

ABC	feature	locally finiteness
ZZZ	factorized	no
OZZ	$2\phi_1$	no
ZZO	$2\phi_1$	no
ZOZ	$3\phi_2$ -like	no
OOZ	factorized	yes
ZOO	factorized	yes
OZO	factorized	no
OOO	$2\phi_1$	yes
XXZ	factorized	no
ZXX	factorized	no
XZX	factorized	no

- For all cases, R^{ABC} are uniquely determined in each disjoint set of recurrence equations specified by appropriate parity conditions.
- We obtained the explicit formulae for them, where their matrix elements are either factorized or expressed as q-hypergeometric series.

- Examples of RLLL relation for ZZZ:

$$R(1 \otimes X \otimes X) = (1 \otimes X \otimes X)R, \quad R(X \otimes X \otimes 1) = (X \otimes X \otimes 1)R,$$

$$-r_1 r_3 R(1 \otimes Z \otimes 1) = (qt_1 t_3 w_1 X \otimes Z \otimes X - r_2 Z \otimes 1 \otimes Z)R,$$

$$R(-qt_1 t_3 w_3 X \otimes Z \otimes X + s_2 Z \otimes 1 \otimes Z) = s_1 s_3 (1 \otimes Z \otimes 1)R,$$

$$t_1 R(X \otimes Z \otimes Z^{-1}(r_3 s_3 - t_3^2 w_3 X^2) + s_2 t_3 Z \otimes 1 \otimes X) = s_3 t_2 (Z \otimes X \otimes 1)R,$$

$$R(t_3 w_3 Z^{-1}(r_1 s_1 - t_1^2 w_1 X^2) \otimes Z \otimes X + s_2 t_1 w_1 X \otimes 1 \otimes Z) = s_1 t_2 w_2 (1 \otimes X \otimes Z)R.$$

$$\pi_Z : X |m\rangle = |m-1\rangle, \quad Z |m\rangle = q^m |m\rangle$$

- Writing down actions of π_Z , we obtain recursion relations for R^{ZZZ} :

$$R_{i,j-1,k-1}^{a,b,c} = R_{i,j,k}^{a,b+1,c+1}, \quad R_{i-1,j-1,k}^{a,b,c} = R_{i,j,k}^{a+1,b+1,c},$$

$$(q^{a+c} r_2 - q^j r_1 r_3) R_{i,j,k}^{a,b,c} = q^{1+b} t_1 t_3 w_1 R_{i,j,k}^{a+1,b,c+1},$$

$$(q^{i+k} s_2 - q^b s_1 s_3) R_{i,j,k}^{a,b,c} = q^{1+j} t_1 t_3 w_3 R_{i-1,j,k-1}^{a,b,c},$$

$$q^j r_3 s_3 t_1 R_{i-1,j,k}^{a,b,c} - q^{j+2} t_1 t_3^2 w_3 R_{i-1,j,k-2}^{a,b,c} + q^{i+k} s_2 t_3 R_{i,j,k-1}^{a,b,c} = q^{a+k} s_3 t_2 R_{i,j,k}^{a,b+1,c},$$

$$q^j r_1 s_1 t_3 w_3 R_{i,j,k-1}^{a,b,c} - q^{j+2} t_1^2 t_3 w_1 w_3 R_{i-2,j,k-1}^{a,b,c} + q^{i+k} s_2 t_1 w_1 R_{i-1,j,k}^{a,b,c} = q^{c+i} s_1 t_2 w_2 R_{i,j,k}^{a,b+1,c}$$

- Fact: Recursion relations for ZZZ consists of 4 disjoint sets, which are specified with the parity pair $(d_1, d_2) = (a + c - j, b - i - k)$.

■ Thm: [Kuniba-Matsuike-Y'22]

- $R^{ZZZ} \in \text{End}(F^{\otimes 3})$ is uniquely determined in each sector and given by

$$R_{i,j,k}^{a,b,c} = \left(\frac{r_2}{t_1 t_3 w_1} \right)^{\frac{d_1}{2}} \left(\frac{s_2}{t_1 t_3 w_3} \right)^{\frac{d_2}{2}} \left(\frac{t_2}{s_1 t_3} \right)^{\frac{d_3}{2}} \left(\frac{t_2 w_2}{s_3 t_1 w_1} \right)^{\frac{d_4}{2}} \\ \times q^\varphi \frac{\Phi_{d_2} \left(\frac{s_1 s_3}{s_2} \right) \Phi_{d_3} \left(\frac{r_3 w_2}{s_3 w_1} \right) \Phi_{d_4} \left(\frac{r_1 w_3}{s_1 w_2} \right)}{\Phi_{-d_1} \left(\frac{q^2 r_1 r_3}{r_2} \right) \Phi_{d_3+d_4} \left(\frac{r_1 r_3 w_3}{s_1 s_3 w_1} \right)}, \quad a, b, c, i, j, k \in \mathbb{Z}$$

$$\varphi = \frac{1}{4} ((d_1 - d_2)(d_1 + d_2 + d_3 + d_4) + d_3 d_4) - d_1,$$

$$\begin{pmatrix} d_1 \\ d_2 \end{pmatrix} = \begin{pmatrix} a + c - j \\ b - i - k \end{pmatrix}, \quad \begin{pmatrix} d_3 \\ d_4 \end{pmatrix} = \begin{pmatrix} -a - b + c + i + j - k \\ a - b - c - i + j + k \end{pmatrix}$$

$$\Phi_m(z) = \frac{1}{(zq^m; q^2)_\infty} \quad (m \in \mathbb{Z}),$$

■ Features:

- The matrix elements of R^{ZZZ} are factorized.
- R^{ZZZ} is *not* locally finite.
- There are 4 sectors specified with the parity pair (d_1, d_2) .

■ Thm: [Kuniba-Matsuike-Y'22]

□ $R^{OZZ} \in \text{End}(F_+ \otimes F \otimes F)$ is uniquely determined and given by

$$R_{i,j,k}^{a,b,c} = \theta(i \geq 0) \left(\frac{r_2}{r_3}\right)^a \left(\frac{s_3}{s_2}\right)^i \left(\frac{t_2 w_2}{\mu s_2}\right)^{-b+j} \left(-\frac{\mu t_3}{r_3}\right)^{-c+k} \frac{(z; q^2)_a}{(q^2; q^2)_a} q^{(a-b+j-1)c - (i-b+j-1)k - aj + bi}$$

$$\times {}_2\phi_1 \left(\begin{matrix} q^{-2i}, z^{-1}q^2 \\ z^{-1}q^{-2a+2} \end{matrix}; q^2, yq^{2i+2j-2a-2b} \right). \quad a, i \in \mathbb{Z}_{\geq 0}, b, c, j, k \in \mathbb{Z}$$

$$\mu = \mu_4, \quad y = \frac{r_3 w_3}{\mu^2 s_3}, \quad z = q^{2k-2c+2} \frac{\mu^2 s_2}{r_2 w_2}$$

■ Features:

- The matrix elements of R^{OZZ} are expressed as q-hypergeometric series.
- R^{OZZ} is *not* locally finite.
- There is only 1 sector.

■ Thm: [Kuniba-Matsuike-Y'22]

- $R^{OOZ} \in \text{End}(F_+ \otimes F_+ \otimes F)$ is uniquely determined and non-trivial iff $\mu_1 / \mu_2 = q^d$ for $d \in \mathbb{Z}$. In that case, it is given by

$$R(d)_{i,j,k}^{a,b,c} = \theta(e \in \mathbb{Z}) \theta(\min(i, j) \geq 0) \delta_{i+j}^{a+b} \quad a, b, i, j \in \mathbb{Z}_{\geq 0}, c, k \in \mathbb{Z}$$

$$\times s_3^i (\mu_2 t_3)^{-a} \left(\frac{\mu_2 s_3}{t_3 w_3} \right)^j \left(\frac{t_3^2 w_3}{r_3 s_3} \right)^e q^{cj-bk} \frac{(q^{2+2e-2j}; q^2)_j (q^{2a+2}; q^2)_{i-a}}{(q^2; q^2)_f (q^{2a-2e}; q^2)_{e-a}}$$

$$e = \frac{1}{2}(a - c + j + k + d), \quad f = \frac{1}{2}(b + c + i - k - d)$$

■ Features:

- The matrix elements of R^{OOZ} are factorized.
- R^{OOZ} is locally finite.
- There is only 1 sector but R^{OOZ} is non-trivial if the parity of $2e$ is even.

- From associativity, we can expect the RRRR equations holds:

$$R_{124}^{ABD} R_{135}^{ACE} R_{236}^{BCF} R_{456}^{DEF} = R_{456}^{DEF} R_{236}^{BCF} R_{135}^{ACE} R_{124}^{ABD}$$

$$A, B, C, D, E, F \in \{X, Z, O\}$$

- Conjecture: [Kuniba-Matsuike-Y'22]

- At least, the following RRRR equations are valid:

$$\begin{aligned} R_{456}^{OOO} R_{236}^{OOO} R_{135}^{ZOO} R_{124}^{ZOO} &= R_{124}^{ZOO} R_{135}^{ZOO} R_{236}^{OOO} R_{456}^{OOO} \\ R_{456}^{ZOO} R_{236}^{OOO} R_{135}^{OOO} R_{124}^{OOZ} &= R_{124}^{OOZ} R_{135}^{OOO} R_{236}^{OOO} R_{456}^{ZOO} \\ R_{456}^{OOZ} R_{236}^{OOZ} R_{135}^{OOO} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OOO} R_{236}^{OOZ} R_{456}^{OOZ} \\ R_{456}^{OOZ} R_{236}^{OOZ} R_{135}^{ZOO} R_{124}^{ZOO} &= R_{124}^{ZOO} R_{135}^{ZOO} R_{236}^{OOZ} R_{456}^{OOZ} \\ R_{456}^{OOO} R_{236}^{ZOO} R_{135}^{OOO} R_{124}^{OZO} &= R_{124}^{OZO} R_{135}^{OOO} R_{236}^{ZOO} R_{456}^{OOO} \\ R_{456}^{OZO} R_{236}^{OOO} R_{135}^{OOZ} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OOZ} R_{236}^{OOO} R_{456}^{OZO} \\ R_{456}^{OOO} R_{236}^{ZOO} R_{135}^{ZOO} R_{124}^{ZZO} &= R_{124}^{ZZO} R_{135}^{ZOO} R_{236}^{ZOO} R_{456}^{OOO} \\ R_{456}^{ZOO} R_{236}^{OOO} R_{135}^{ZOO} R_{124}^{ZOZ} &= R_{124}^{ZOZ} R_{135}^{ZOO} R_{236}^{OOO} R_{456}^{ZOO} \\ R_{456}^{ZOO} R_{236}^{ZOO} R_{135}^{OOO} R_{124}^{OZZ} &= R_{124}^{OZZ} R_{135}^{OOO} R_{236}^{ZOO} R_{456}^{ZOO} \\ R_{456}^{ZZO} R_{236}^{OOO} R_{135}^{OOZ} R_{124}^{OOZ} &= R_{124}^{OOZ} R_{135}^{OOZ} R_{236}^{OOO} R_{456}^{ZZO} \\ R_{456}^{ZOZ} R_{236}^{OOZ} R_{135}^{OOO} R_{124}^{OOZ} &= R_{124}^{OOZ} R_{135}^{OOO} R_{236}^{OOZ} R_{456}^{ZOZ} \\ R_{456}^{OZZ} R_{236}^{OOZ} R_{135}^{OOZ} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OOZ} R_{236}^{OOZ} R_{456}^{OZZ} \\ R_{456}^{ZOO} R_{236}^{ZOO} R_{135}^{ZOO} R_{124}^{ZZZ} &= R_{124}^{ZZZ} R_{135}^{ZOO} R_{236}^{ZOO} R_{456}^{ZOO} \\ R_{456}^{ZZZ} R_{236}^{OOZ} R_{135}^{OOZ} R_{124}^{OOZ} &= R_{124}^{OOZ} R_{135}^{OOZ} R_{236}^{OOZ} R_{456}^{ZZZ} \end{aligned}$$

$$\begin{aligned} R_{456}^{OOO} R_{236}^{OZO} R_{135}^{OZO} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OZO} R_{236}^{OZO} R_{456}^{OOO} \\ R_{456}^{OOO} R_{236}^{OZO} R_{135}^{ZZO} R_{124}^{ZOO} &= R_{124}^{ZOO} R_{135}^{ZZO} R_{236}^{OZO} R_{456}^{OOO} \\ R_{456}^{OZO} R_{236}^{OOO} R_{135}^{ZOZ} R_{124}^{ZOO} &= R_{124}^{ZOO} R_{135}^{ZOZ} R_{236}^{OOO} R_{456}^{OZO} \\ R_{456}^{OOO} R_{236}^{ZZO} R_{135}^{OZO} R_{124}^{OZO} &= R_{124}^{OZO} R_{135}^{OZO} R_{236}^{ZZO} R_{456}^{OOO} \\ R_{456}^{OOZ} R_{236}^{ZOZ} R_{135}^{OOO} R_{124}^{OZO} &= R_{124}^{OZO} R_{135}^{OOO} R_{236}^{ZOZ} R_{456}^{OOZ} \\ R_{456}^{ZOO} R_{236}^{OZO} R_{135}^{OZO} R_{124}^{OOZ} &= R_{124}^{OOZ} R_{135}^{OZO} R_{236}^{OZO} R_{456}^{ZOO} \\ R_{456}^{OZO} R_{236}^{OZO} R_{135}^{OZZ} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OZZ} R_{236}^{OZO} R_{456}^{OZO} \\ R_{456}^{OOZ} R_{236}^{OZZ} R_{135}^{OZO} R_{124}^{OOO} &= R_{124}^{OOO} R_{135}^{OZO} R_{236}^{OZZ} R_{456}^{OOZ} \\ R_{456}^{OZO} R_{236}^{OZO} R_{135}^{ZZZ} R_{124}^{ZOO} &= R_{124}^{ZOO} R_{135}^{ZZZ} R_{236}^{OZO} R_{456}^{OZO} \\ R_{456}^{OOZ} R_{236}^{ZZZ} R_{135}^{OZO} R_{124}^{OZO} &= R_{124}^{OZO} R_{135}^{OZO} R_{236}^{ZZZ} R_{456}^{OOZ} \end{aligned}$$

Remark: Each equation is checked for over 10000 outer lines by computer.

■ Proposition: [Kuniba-Matsuike-Y'22]

- $R^{ZZZ} \in \text{End}(F^{\otimes 3})$ satisfies the following intertwining relation of the quantum coordinate ring $A_q(A_2)$:

$$R^{ZZZ} \circ (\pi_1 \otimes \pi_2 \otimes \pi_1(\Delta^{\text{op}}(g))) = (\pi_2 \otimes \pi_1 \otimes \pi_2(\Delta(g))) \circ R^{ZZZ} \quad \forall g \in A_q(A_2)$$

- $\pi_i = \pi_Z \circ \varrho_i$, where ϱ_1 and ϱ_2 are respectively given by t_{ij} : generators of $A_q(A_2)$

$$\begin{pmatrix} t_{11} & t_{12} & t_{13} \\ t_{21} & t_{22} & t_{23} \\ t_{31} & t_{32} & t_{33} \end{pmatrix} \mapsto \begin{pmatrix} Z^{-1}(u_1 - g_1 h_1 X^2) & g_1 X & 0 \\ -q h_1 X & Z & 0 \\ 0 & 0 & u_1^{-1} \end{pmatrix}, \begin{pmatrix} u_2^{-1} & 0 & 0 \\ 0 & Z^{-1}(u_2 - g_2 h_2 X^2) & g_2 X \\ 0 & -q h_2 X & Z \end{pmatrix}$$

- π_i s are **not** irreducible. $\pi_Z : X |m\rangle = |m-1\rangle, \quad Z |m\rangle = q^m |m\rangle$

- Identification of parameters is done as follows:

$$u_1 = u_2 (= : u) \qquad g_1 h_1 = g_2 h_2 (= : p)$$

$$\frac{r_1}{t_1} = \frac{r_2}{t_2}, \quad \frac{s_2}{t_2} = \frac{s_3}{t_3}, \quad \frac{r_2}{r_1 r_3} = u, \quad \frac{s_1 s_3}{s_2} = u^2, \quad \frac{t_1^2 w_1}{r_1 s_1} = \frac{t_2^2 w_2}{r_2 s_2} = \frac{t_3^2 w_3}{r_3 s_3} = \frac{p}{u}.$$

■ Summary:

- We considered three kinds of L -operators L^X, L^Z, L^O and $RLLL$ relations which they satisfy. They can be regarded as q -Oscillator or q -Weyl algebra valued six vertex models.
- We solved these $RLLL$ relations and obtained explicit formulae for R^{ABC} .
- For all cases, R^{ABC} are uniquely determined in each disjoint set of recurrence equations specified by appropriate parity conditions.
- Their matrix elements are either factorized or expressed as q -hypergeometric series.
- By computer experiments, we conjectured $RRRR$ equations for R^{ABC} .
- We found R^{ZZZ} satisfies an intertwining relation for *reducible* representations of $A_q(A_2)$.

■ Outlook:

- Root of unity, Bazhanov-Baxter model: F, F_+ become finite dimensional
- Boundary integrability in 3D: $K(LGLG) = (GLGL)K$

- Reduction to 2D, generalized chiral Potts model

- Characterization in terms of PBW bases of U_q^+

$$e_2^{(a)} e_{12}^{(b)} e_1^{(c)} = \sum_{i,j,k \in \mathbb{Z}_{\geq 0}} (R^{OOO})_{i,j,k}^{a,b,c} e_1^{(k)} e_{21}^{(j)} e_2^{(i)}$$